

$\begin{array}{c} \textbf{MANAGEMENT OF THERAPEUTIC ANTICOAGULATION IN PREGNANCY, DELIVERY AND} \\ \textbf{POSTPARTUM} \end{array}$

Pat	ient Information			in event:	current	1 0 3
			PE Gestatio	in on at time of e	current vent:	pregnancy
Thrombosis	History:	Or/	Long Anticoag		icoagulation	pre pregnancy
		Indication	on:			
Anticoagula	tion during pregnancy					
Agent:						
Dose:						
Planned Dur	ation of Treatment:					
Delivery Pla	ın					
□ Spontan	eous labour (ensure patien	t knows not	to take Ll	MWH if susp	ects/in labour)	
Or/ □ Planned	delivery: Induction of	of labour	ca	aesarean secti	on	
Date of plans	ned delivery:					
Last dose of (Therapeutic	LMWH: Date LWMH should be discont				Timesarean section)	
Authorized	by Dr			Date		······
Post Deliver	y					
1. Delivery	y with no bleeding compl	ications:				
a.	Prophylactic dose LWM hours post epidural cather				4 hour	s post delivery or 4
b.	Treatment dose LMWH dose (use postpartum we					5mg/kg single daily
Authorized b	y Dr	I	Date			

2. Delivery with bleeding complications:

Options are to delay dose of LMWH, o should be discussed with the on-call ha	or in rare situations consider the use of UF Hepari nematology consultant	in. These cases
Plan:		
Duration of postnatal anticoagulation	n	
anticoagulation at treatment dose and a treatment dose anticoagulation will be	rrent pregnancy require a minimum of 6 weeks post a total duration of therapy of 3 months (therefore extended beyond 6 weeks postpartum depending from LMWH to Warfarin or a DOAC until Day 5 eeding women	for some women on when the VTE
Planned duration of PN anticoagulat	tion	
Options: Continue LWMH	☐ LMWH/Warfarin	□ DOAC
(Refer to GG&C Guideline Thromboe Management for dosing regimens)	embolic Disease in Pregnancy and the Puerperiu	ım – Acute
Postnatal Follow-up Arrangements:		
Authors: Dr C Bagot, Dr J Brennand	d, Dr V Brace	
Approved by Obstetric Guideline Gr Governance Group:	roup:Appro	oved by Obstetric
Implementation Date:04/12/18 Date:01/12/23		