This information was up to date at the time of release to the Heads of Midwifery.

The editorial board does not accept liability for any errors or omissions following its subsequent publication.

Updating arrangements for the formulary should be decided upon and implemented at a local level.

Miconazole 2% Cream (Daktarin®)	
Legal status (GSL, P or POM on exemption list, or PGD)	P - midwife may supply as medicine is on midwives exemptions list
Patient group	Neonates with suspected fungal nappy rash
Clinical indication	Fungal skin infections
Pharmacology (Onset and duration of action where appropriate)	An antifungal against common dermatophytes and yeasts and also has antibacterial activity against certain gram-positive bacilli and cocci.
	Inhibits the biosynsthesis of ergosterol in fungi and changes the composition of other lipid components in the membrane, resulting in fungal cell necrosis.
	Symptoms such as pruritus respond rapidly.
Pharmaceutical form, strength, route of administration	Cream containing 2% miconazole in 15g and 30g tubes. For topical application.
Dose, frequency and maximum number of doses or period of time for administration or supply	Apply twice a day continuing for 10 days after lesions have healed
Contra-indications/exclusion criteria	 known hypersensitivity to any component of the medicine or to other imidazole derivatives
Cautions and action that will be taken if a caution applies	 discontinue if local sensitisation or allergic reaction occurs contains benzoic acid and butylated hydroxyanisole which may cause local skin reactions (e.g. contact dermatitis), or irritation to the eyes and mucous membranes. check for and document any allergies if a caution applies consult with a doctor document in neonatal record

Miconazole 2% Cream (Daktarin®) Medicine interactions and action other interactions unlikely due to limited systemic absorption that will be taken if a patient is . if there is a clinically significant drug interaction suspected, taking a medicine that may consult with a doctor before administration or supply interact document consultation in neonatal record refer to current BNF for latest information on interactions Potential adverse reactions and occasional local irritation side effects including actions to • other reaction reported: angioedema be taken if adverse drug reaction is suspected if a serious adverse reaction is suspected please report to the MHRA Yellow Card Scheme. http://yellowcard.mhra.gov.uk/ **Overdose** no problem expected • Excessive use can result in skin irritation. Usually resolves on discontinuation For cutaneous use only. If accidental ingestion occurs seek urgent medical review for further advice contact National Poisons Centre 0344 892 0111 **Action if patient declines** refer to authorised prescriber or doctor document in neonatal record Additional advice and If neonate being treated for oral thrush dual therapy with topical information miconazole in napkin area should be commenced

Miconazole 2% Cream (Daktarin®)	
Patient monitoring arrangements during and after treatment and follow-up required	Refer to neonatal doctor if worsening symptoms
Particular storage requirements	-

References

- 1. Summary of Product Characteristics Daktarin 2% Cream. Text revision 08.08.2016. Accessed 22.01.2020 http://www.medicines.org.uk
- 2. http://www.bnf.org