

**NHS Borders
Management of Upper GI Bleed**

**Proton-Pump-Inhibitor (PPI) Infusion
Chart
(Also prescribe on the medicine
chart)**

Ward/Hosp..... Cons.....

CHI

--	--	--	--	--	--	--	--	--	--

Surname.....

First Name.....

Address.....

USE BLOCK LETTERS OR PATIENT LABEL

Date of Infusion ___ / ___ / 20__			* Please ensure patient receives continuous infusion for 70 hours post-endoscopy when indicated. Infusion to commence immediately after loading dose				Prescribed by (Doctors Signature)	
Bottle No.	FLUID		Rate of Flow per Hour ml/hr	Batch No.	ADDITIONS	Use 24hr Clock		Administered by (Signature)
	TYPE	Amount ml				Start	Finish	
LD	Sodium Chloride 0.9%	100ml	200ml/hr		Omeprazole 80mg			
1	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
2	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
3	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
4	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
5	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
6	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			
7	Sodium Chloride 0.9%	100ml	10ml/hr		Omeprazole 80mg			

Blatchford Score Guidance (On Admission)				Patients Score =		
Blood urea (mmol/L)	Score	Haemoglobin (g/L) for women	Score	Other markers	Score	
6.5 – 7.9	2	10.0 – 12.0	1	Pulse >100 (per min)	1	
8.0 - 9.9	3	<10.0	6	Presentation with malaena	1	
10.0 – 25	4			Presentation with syncope	2	
> 25	6	Systolic BP (mm Hg)		Hepatic disease	2	
Haemoglobin (g/L) for men		100–109	1	Cardiac failure	2	
12.1 - 13.0	1	90–99	2			
10.1 – 12.0	3	<90	3			
<10.0	6					

Blatchford Score 0 – 1 (Low Risk of Bleeding) Blatchford Score 2 – 5 (Intermediate Risk of Bleeding)
Blatchford Score > 6 (High Risk of Bleeding)

Date and Time of Endoscopy: _____

Endoscopic Findings: _____

Predicted Date and Time to Stop Infusion: _____

