NGU treatment flowchart

Microscopically proven NGU

1st line: Doxycycline 100mg BD 7 days
(2nd line: Azithromycin 1g D1, 500mg D2, 500mg D3)

Abstain for 2 weeks from start of treatment and until symptoms have resolved.
Treat regular partners with same regime and test for Ct/GC

Microscopically proven persistent/recurrent non-gonococcal urethritis
Confirm adherence to previous treatment
Confirm partner treated and not re-infection

Send *M.genitalium* test

Empirical treatment appropriate (eg severe symptoms, likely to have sexual contact)

Has completed 1 week of doxycycline within 2 weeks?

YES:

Azithromycin 1g D1, 500mg D2, 500mg D3
+/- metronidazole 400mg BD 5 days

Consider treating regular partners with same regimen

Await *M.genitalium* result

NO:

Doxycycline 100mg BD 1 week

Await *M.genitalium* result

Appropriate to wait for result (eg mild symptoms, unlikely to have sexual contact)

M.genitalium NEGATIVE:

Azithromycin 1g D1, 500mg D2, 500mg D3
+metronidazole 400mg BD 5 days

Consider treating regular partners with same

If persists or recurs consider other causes and refer Urology

M.genitalium POSITIVE:
Follow M.gen protocol

Hea cols. 10th Edition.