



## CLINICAL GUIDELINE

# Hospital Infection Management Guidelines Empirical Antibiotic Therapy in Adults

A guideline is intended to assist healthcare professionals in the choice of disease-specific treatments.

Clinical judgement should be exercised on the applicability of any guideline, influenced by individual patient characteristics. Clinicians should be mindful of the potential for harmful polypharmacy and increased susceptibility to adverse drug reactions in patients with multiple morbidities or frailty.

If, after discussion with the patient or carer, there are good reasons for not following a guideline, it is good practice to record these and communicate them to others involved in the care of the patient.

<b>Version Number:</b>	12
<b>Does this version include changes to clinical advice:</b>	Yes
<b>Date Approved:</b>	18 <sup>th</sup> February 2025
<b>Date of Next Review:</b>	31 <sup>st</sup> August 2026
<b>Lead Author:</b>	Scott Gillen
<b>Approval Group:</b>	Antimicrobial Utilisation Committee

### Important Note:

The online version of this document is the only version that is maintained. Any printed copies should therefore be viewed as 'Uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

# Hospital Infection Management Guidelines Empirical Antibiotic Therapy in Adults

**STOP AND THINK BEFORE ANTIBIOTIC THERAPY:** 1 in 5 antimicrobial courses associated with adverse events including *C.difficile*, drug interactions/ toxicity, device related infections and *S. aureus* bacteraemia. **THINK SEPSIS** if NEWS ≥ 5. **Send 2 blood culture sets (4 bottles in total), ensuring each bottle is filled with 10ml of blood before starting antibiotics.**

**RECORD** antimicrobial indication and duration on HEPMA

**REVIEW** patient and results. **RECORD clinical response and prescription daily.** **Can you SIMPLIFY, SWITCH or STOP?** If Clinical improvement + eating/drinking + deep seated/complex infection not suspected then **IVOST** ( See [IVOST Guidelines](#) )

and **RECORD** duration of remaining oral therapy. **RECORD the STOP date for oral antimicrobial on HEPMA**

**REVIEW** all IV antimicrobial and prescription **DAILY** and **RECORD** duration /review date. **INFORM** patient of reason for antimicrobial and likely duration.

**NB** Doses recommended based on normal renal/liver function - see BNF or Renal handbook for dosing advice. For info on antimicrobial contra-indications, cautions and monitoring see BNF.

**Definition of SEPSIS:** **INFECTION** (includes Systemic Inflammatory Response Syndrome (SIRS\*)) **WITH** evidence of **ORGAN HYPOPERFUSION** (≥ 2 of: Confusion, < 15 GCS or Resp Rate ≥ 22/ min or Systolic BP ≤ 100 mm Hg).

**Ensure SEPSIS 6 within one hour:** 1. Blood cultures (& any other relevant samples), 2. IV Antibiotic administration, 3. Oxygen to maintain target saturation, 4. Measure lactate, 5. IV fluids, 6. Monitor urine output hourly.

\*SIRS indicated by Temp < 36°C or > 38°C, HR > 90 bpm, RR > 20/ min & WCC < 4 or > 12 x10<sup>9</sup>/ L. SIRS is not specific to bacterial infection (also viral & non-infective causes).

 <h3>Lower Respiratory Tract Infections</h3> <p><b>Infective Exacerbation COPD</b> Antibiotics only if purulent sputum (send for culture along with viral gargle) <b>Dual antibiotic therapy not recommended &amp; increases risk of harm</b> Oral + Doxycycline 200mg as a one-off single dose then 100mg daily or Oral Amoxicillin 500mg 8 hrly or Oral + Clarithromycin 500mg 12 hrly <b>Duration 5 days</b></p> <p><b>Suspected Viral Respiratory Tract Infection</b> Antibiotics NOT required unless secondary bacterial infections e.g. COPD exacerbation with purulent sputum (see above) If consolidation treat as per CAP below <a href="#">COVID-19 guidelines</a> <a href="#">Flu guidelines</a></p> <p><b>Uncertain if LRTI/ UTI</b> Send MSSU, sputum and viral gargle Oral + Co-trimoxazole 960mg 12 hrly or Oral + Doxycycline 100mg 12 hrly Do NOT prescribe Co-amoxiclav <b>Review/ clarify diagnosis at 48 hours</b> <b>Duration</b> if diagnosis remains uncertain <b>MAXIMUM 5 days</b></p>	 <h3>Skin/ Soft Tissue Infections</h3> <p><b>Mild skin/soft tissue infection</b> Oral Flucloxacillin 1g 6 hrly or if true penicillin/beta-lactam allergy Oral + Co-trimoxazole 960mg 12 hrly or Oral + Doxycycline 100mg 12 hrly <b>Duration 5 days</b></p> <p><b>Moderate / Severe Cellulitis</b> Consider OPAT/ ambulatory care (consult local management pathway). If requires inpatient management: IV Flucloxacillin 2g 6 hrly If <b>MRSA suspected or if true penicillin/ beta-lactam allergy</b> IV Vancomycin** If <b>rapidly progressive</b> Add IV Clindamycin 600mg 6 hrly Consider <b>CDI risk</b> <b>Duration 7-10 days</b> (IV/oral)</p>	 <h3>Gastrointestinal Infections</h3> <p><b>Gastroenteritis</b> Confirm travel history/other risk factors <b>Antibiotics not usually required</b> and may be deleterious in <i>E.coli</i> O157 Consider viral causes</p> <p><b>C. difficile infection (CDI)</b> See <a href="#">CDI Guidelines</a> Treat before lab confirmation if high clinical suspicion. Discontinue if toxin negative</p>	 <h3>Urinary Tract Infections</h3> <p><b>UTI in Pregnancy</b> See NHS GGC Obstetric guidance</p> <p><b>Lower UTI/cystitis</b> Don't treat asymptomatic bacteriuria. Obtain urine culture prior to antibiotic. In women often self-limiting, consider delayed prescribing. Antibiotics if significant symptoms ≥ 2 of dysuria, frequency, urgency, nocturia, haematuria, (and for adult women &lt; 65 years +ve urine nitrite) Oral Nitrofurantoin 50mg 6 hourly or Nitrofurantoin 100mg MR 12 hourly or Oral + Trimethoprim 200mg 12 hrly <b>Duration: Females 3 days, Males 7 days</b> If eGFR &lt; 30 mL/min/1.73 m<sup>2</sup> Nitrofurantoin contraindicated, Trimethoprim use with caution</p> <p><b>Upper UTI</b> Obtain urine for culture prior to antibiotic. Exclude pneumonia if loin/back pain <b>Non-severe/without sepsis</b> Oral + Ciprofloxacin 500mg 12 hrly Or Oral + Co-trimoxazole 960 mg 12 hrly if trimethoprim sensitive organism. <b>Duration 7 days</b></p>	 <h3>Bone/ Joint Infections</h3> <p><b>Septic arthritis/Osteomyelitis / Prosthetic joint infection</b> Urgent orthopaedic referral if underlying metal work or recent surgery. Obtain blood cultures (and if not acutely unwell/ septic, obtain synovial/ other deep samples) prior to antibiotic therapy</p> <p><b>Native joint</b> IV Flucloxacillin 2g 6 hrly If <b>MRSA suspected or if true penicillin/beta-lactam allergy</b> IV Vancomycin** If considered high risk for Gram negative infection e.g. immunocompromised, recurrent UTI or sickle cell disease: <b>ADD</b> IV Gentamicin**Δ (max 4 days) <b>Duration and IVOST:</b> discuss with Infection Specialist at 72 hours. Usually 4-6 weeks (IV/oral) if diagnosis confirmed. <b>Prosthetic joint</b> Antibiotic therapy should not be started in a clinically stable patient until intra-operative samples obtained IV Vancomycin** + IV Gentamicin**Δ (max 4 days) <b>Duration and IVOST: discuss with Infection Specialist at 72 hours</b></p>	 <h3>CNS Infections</h3> <p><b>LP safe without CT scan UNLESS:</b> seizures, GCS ≤ 12, CNS signs, papilloedema or immunosuppression. If CT: Blood cultures and antibiotics BEFORE CT scan. Use Meningitis/ Encephalitis order set on Trakcare, Blood and CSF Glucose. <b>LP contraindicated if:</b> Brain shift, rapid GCS reduction, Resp/ cardiac compromise, severe sepsis, rapidly evolving rash, infection at LP site, coagulopathy, thrombocytopenia, anticoagulant drugs</p> <p><b>Possible bacterial meningitis</b> IV Ceftriaxone 2g 12 hrly or if previous penicillin anaphylaxis IV Chloramphenicol 25mg/kg (max 2g) 6 hrly If bacterial meningitis strongly suspected: <b>ADD</b> IV Dexamethasone 10mg 6 hrly (for 4 days) Prior to, or at the same time as antibiotics and refer to ID If age ≥ 60 years, immunosuppressed, pregnant, alcohol excess, liver disease or if listeria meningitis suspected: <b>ADD</b> IV Amoxicillin 2g 4 hrly to Ceftriaxone <b>Duration of antibiotics: Discuss with Infection Specialist</b></p>	 <h3>Severe Systemic Infection Source Unknown</h3> <p><b>Sepsis where source unknown</b> <b>Review all anatomical systems,</b> perform CXR and consider other imaging/ laboratory investigations Review previous microbiology results and discuss with an infection specialist if previous gentamicin resistance <b>Review diagnosis DAILY</b> <b>Add cover for S aureus infection if:</b> healthcare associated, recent hospitalisation, post-op wound/ line related, PWID <b>Add cover for MRSA infection if;</b> recent MRSA carrier or previous infection <b>Add cover for Streptococcal infection if;</b> pharyngitis/erythroderma/hypotension</p> <p><b>Source unknown</b> IV Amoxicillin 1g 8 hrly + IV Gentamicin**Δ (max 4 days) If <b>S aureus suspected</b> <b>ADD</b> IV Flucloxacillin 2g 6 hrly If <b>MRSA suspected or if true penicillin/ beta-lactam allergy</b> IV Vancomycin** + IV Gentamicin**Δ (max 4 days) If <b>severe Streptococcal infection suspected</b> <b>ADD</b> IV Clindamycin 600mg 6 hrly If eGFR &lt; 20mL/min/1.73 m<sup>2</sup>, <b>REPLACE</b> Gentamicin with Oral/IV + Ciprofloxacin <b>Duration: Review with response/ micro results at 72 hours</b></p>	 <h3>Immunocompromised Patient</h3> <p><b>Immunocompromised Patient</b> Recent Chemotherapy (&lt; 3 weeks), high dose steroids (e.g. prednisolone &gt; 20mg/day for &gt; 2 weeks), other immunosuppressants (e.g. anti-TNF, cyclophosphamide), Stem cell/solid organ transplant or primary immunodeficiency</p> <p><b>Neutropenic Sepsis</b> Neutrophils ≤ 0.5 x 10<sup>9</sup>/ L + fever (temperature &gt; 38°C or 37.5°C on 2 occasions 30 min apart) / hypothermia &lt; 36°C OR chills, shivers, sweats or other symptoms suggestive of infection. All patients who have received recent chemotherapy and who exhibit any of the symptoms above are presumed to be neutropenic and septic.</p> <p><b>Immunocompromised with fever BUT normal neutrophils AND source of infection identified</b> Manage as per infection management guidelines based on anatomical source.</p> <p><b>Neutropenic sepsis or Immunocompromised with fever and source of infection unknown;</b> See guideline Initial Management of Neutropenic Sepsis or Sepsis of Unknown Source in Immunocompromised Adults which is available on StaffNet by clicking: →Clinical Info →NHSGGC Clinical Guideline Platform →Adult infection Management →Secondary Care - Treatment <a href="#">Neutropenic Sepsis or Sepsis of unknown source in immunocompromised adults (168)   Right Decisions</a></p>																
Urgent Blood Cultures then IV Antimicrobial Therapy within ONE hour																							
<h3>Community Acquired Pneumonia (CAP)</h3> <p>Assess for SEPSIS Calculate CURB 65 score: • Confusion (new onset) • Urea &gt; 7 mmol/L • RR ≥ 30 breaths/ min • BP – diastolic ≤ 60 mmHg or systolic &lt; 90 mmHg • Age ≥ 65 years If patient admitted from a care home treat as CAP. If severe, ensure atypical screen sent.</p> <p><b>Non-severe CAP</b> CURB65 score: ≤ 2 (and no sepsis) Oral Amoxicillin 500mg 8 hrly or Oral + Doxycycline 200mg as a one-off single dose then 100mg daily or Oral + Clarithromycin 500mg 12 hrly <b>Duration 5 days</b></p> <p><b>Severe CAP</b> <b>CURB 65 score ≥ 3 or CAP (with any CURB 65 score) PLUS sepsis :</b> Oral + Clarithromycin 500mg 12 hrly <b>PLUS either:</b> IV Amoxicillin 1g 8 hrly or if requiring HDU/ ICU level care IV Co-amoxiclav 1.2g 8 hrly If true penicillin/beta-lactam allergy or Legionella strongly suspected Oral + Levofloxacin <b>Monotherapy</b> 500mg 12 hrly (NB oral bioavailability 99 – 100%) <b>Duration 5 days</b> (IV/oral) Legionella 10-14 days</p>	<h3>Hospital Acquired Pneumonia (HAP)</h3> <p>Diagnosis of HAP is difficult and it is often over-diagnosed. Consider other causes of clinical deterioration including hospital onset COVID-19 and review diagnosis early. Seek senior advice. Assess severity based on CURB 65 score. If within 4 days of admission or admitted from care home Treat as CAP If ≤ 7 days post hospital discharge or ≥ 5 days after admission: <b>Non-severe HAP</b> Oral therapy recommended Oral + Doxycycline 100mg 12 hrly or Oral + Co-trimoxazole 960mg 12 hrly <b>Duration 5 days</b> Trimethoprim use with caution may ↑ K<sup>+</sup> and decrease renal function. Monitor <b>Severe HAP</b> IV Co-amoxiclav 1.2g 8 hourly + IV Gentamicin**Δ (max 4 days) or if true penicillin/beta-lactam allergy Oral + Levofloxacin 500mg 12 hrly monotherapy <b>Duration 5 days</b> (IV/oral) If critically ill discuss with Infection Specialist</p>	<h3>Suspected Necrotising Fasciitis</h3> <p>Consider in SSTI with disproportionate pain or presence of acute organ dysfunction/ hypoperfusion including hypotension. <b>Seek urgent surgical/ orthopaedic review. Urgent DEBRIDEMENT/ EXPLORATION may be required</b> IV Flucloxacillin 2g 6 hrly + IV Benzylpenicillin 2.4g 6 hrly + IV Metronidazole 500mg 8 hrly + IV Clindamycin 1.2g 6 hrly + IV Gentamicin**Δ (max 4 days) If <b>MRSA suspected or if true penicillin/ beta-lactam allergy</b> <b>REPLACE</b> Flucloxacillin + Benzylpenicillin with IV Vancomycin** <b>Rationalise therapy within 48-72 hours</b> Based on: response, microbiology results infection specialist review <b>Duration 10 days</b> (IV/oral) or as per infection specialist</p>	<h3>Intra-abdominal sepsis</h3> <p>IV Amoxicillin 1g 8 hrly <b>+Oral/ IV Metronidazole 400mg / 500mg 8 hrly</b> <b>+IV Gentamicin**Δ (max 4 days) )</b> If eGFR &lt; 20 mL/min/1.73 m<sup>2</sup> IV Piperacillin/Tazobactam 4.5g 12 hourly (Monotherapy) If true penicillin/beta-lactam allergy IV Vancomycin ** <b>+Oral/ IV Metronidazole 400/ 500mg 8 hrly</b> <b>+IV Gentamicin**Δ (max 4 days)</b> If eGFR &lt; 20mL/min/1.73 m<sup>2</sup> <b>** IV/Oral Ciprofloxacin</b> <b>+Oral/ IV Metronidazole 400/ 500mg 8 hrly</b> <b>Total Duration 5 days</b> (IV/oral) Assuming source control See <a href="#">Advice for Antibiotic therapy following 4 days IV gentamicin</a></p> <p><b>Biliary tract infection</b> As above except metronidazole not routinely required unless severe</p> <p><b>Pancreatitis</b> Does not require antibiotic therapy unless complicated by cholangitis.</p>	<h3>Spontaneous Bacterial Peritonitis (SBP)</h3> <p>SBP confirmed if ascitic counts Manual : WCC &gt;500/mm<sup>3</sup> or neutrophils &gt;250/mm<sup>3</sup> or EDTA automated count: WCC &gt;0.5 or polymorphs &gt;0.25 x10<sup>9</sup>/L See <a href="#">Cirrhosis bundle</a> If not receiving co-trimoxazole prophylaxis: Oral + Co-trimoxazole 960mg 12 hourly If receiving co-trimoxazole prophylaxis: IV Piperacillin/Tazobactam 4.5g 8 hourly or if true penicillin/beta-lactam allergy Oral + Levofloxacin 500mg 12 hrly <b>Duration 7 days</b> (IV/oral)</p> <p><b>Decompensated Chronic liver Disease with Sepsis Unknown Source</b> IV Piperacillin/Tazobactam 4.5g 8 hourly or if true penicillin/beta-lactam allergy Oral + Levofloxacin 500mg 12 hrly <b>Duration 7 days</b> (IV/oral)</p>	<h3>Catheter related UTI</h3> <p>Remove/ replace catheter and send urine for culture. Don't treat asymptomatic bacteriuria</p> <p><b>Symptomatic bacteriuria without sepsis</b> Give single dose of IV Gentamicin**Δ immediately prior to catheter removal or if IV route not available give single dose of oral + Ciprofloxacin 500mg 30 minutes before catheter change. If eGFR &lt; 20 mL/min/1.73 m<sup>2</sup> <b>** Ciprofloxacin 500mg single dose</b> <b>Symptomatic bacteriuria with sepsis</b> Treat as per pyelonephritis/ culture results. <b>Duration 7 days</b> (IV/oral)</p>	<h3>Diabetic foot infection/ osteomyelitis</h3> <p>Assess ulcer size, probes to bone, neuropathy, peripheral vascular disease, MRSA risk. For outpatient therapy consult diabetic clinic guidelines IV Flucloxacillin 2g 6 hrly + Oral Metronidazole 400mg 8 hrly If SEPSIS or SIRS ≥ 2 <b>Add IV Gentamicin**Δ (max 4 days)</b> If <b>MRSA suspected or if true penicillin/beta-lactam allergy</b> IV Vancomycin** + Oral Metronidazole 400mg 8hrly (Metronidazole oral bioavailability 80-100%) <b>If SEPSIS or SIRS ≥ 2:</b> <b>Add IV Gentamicin**Δ (max 4 days)</b> If eGFR &lt; 20 mL/min/1.73 m<sup>2</sup> <b>REPLACE</b> Gentamicin with Oral + Ciprofloxacin <b>Duration/IVOST</b> <b>Discuss with Infection Specialist</b></p>	<h3>Possible infective Endocarditis</h3> <p>Always seek senior specialist advice and refer to cardiology. Native heart valve IV Amoxicillin 2g 4 hrly + IV Flucloxacillin 2g 6 hrly if &lt; 85kg (4 hrly if ≥ 85kg) + IV Gentamicin Δ (synergistic dosing) If <b>MRSA/ resistant organisms suspected or if true penicillin/beta-lactam allergy</b> IV Vancomycin** + IV Gentamicin Δ (synergistic dosing) <b>Prosthetic heart valve</b> IV Vancomycin** + IV Gentamicin Δ (synergistic dosing) <b>Discuss with Infection Specialist within 72 hours</b> *See Synergistic Gentamicin for Endocarditis in Adults guideline on StaffNet for dosing</p>	<p><b>Patients with Stem Cell Transplant or receiving chemotherapy for Acute Leukaemia</b> NEWS ≤ 6 See High Risk treatment above. NEWS ≥ 7 Critical Risk See Neutropenic Sepsis guidelines (see above for pathway to this on StaffNet)</p>															
<h3>Pneumonia</h3>																							
<p><b>**Gentamicin/ **Vancomycin</b> Gentamicin / Vancomycin adult dosing calculators are available via 'Clinical Info' icon on staff intranet/ GGC Medicines App. See GGC Therapeutics Handbook for Prescribing advice. Use GGC Prescribing, Administration, Monitoring charts. Vancomycin If creatinine not available give Vancomycin loading dose as per actual body weight Gentamicin Δ Avoid Gentamicin in decompensated liver disease or myasthenia gravis, or known family history of aminoglycoside auditory toxicity or maternal relative with deafness due to mitochondrial mutation A1555G</p>	<p>If creatinine not available give gentamicin as follows:</p> <table border="1"> <thead> <tr> <th>Actual Body Weight</th> <th>Gentamicin Dose</th> <th>Actual Body Weight</th> <th>Gentamicin Dose</th> </tr> </thead> <tbody> <tr> <td>&lt; 40 kg</td> <td>5 mg/kg</td> <td>60 - 69 kg</td> <td>320 mg</td> </tr> <tr> <td>40 - 49 kg</td> <td>240 mg</td> <td>70 - 79 kg</td> <td>360 mg</td> </tr> <tr> <td>50 - 59 kg</td> <td>280 mg</td> <td>≥ 80 kg</td> <td>400 mg</td> </tr> </tbody> </table> <p>NB If CKD5 give 2.5 mg/kg (max 180 mg)</p>	Actual Body Weight	Gentamicin Dose	Actual Body Weight	Gentamicin Dose	< 40 kg	5 mg/kg	60 - 69 kg	320 mg	40 - 49 kg	240 mg	70 - 79 kg	360 mg	50 - 59 kg	280 mg	≥ 80 kg	400 mg	<h2>!! Important Antibiotic Drug Interactions &amp; Safety Information !!</h2> <ul style="list-style-type: none"> <li><b>+Doxycycline/ Quinolone:</b> reduced absorption with iron, calcium, magnesium &amp; some nutritional supplements. See BNF (Appendix1) or see pharmacy for advice.</li> <li><b>+Clarithromycin/ Quinolone:</b> risk of serious drug interactions see BNF (appendix 1) or seek pharmacy advice. May also prolong the QTc interval, avoid (where possible) if other QTc risk factors. If oral route compromised give IV (see BNF for dose).</li> <li><b>+ Quinolones</b> e.g. Ciprofloxacin, Levofloxacin Stop treatment at first signs of a serious adverse reaction (e.g. tendonitis), prescribe with caution for people over 60 years and avoid co administration with a corticosteroid. See BNF for dosing advice in reduced renal function.</li> <li><b>Trimethoprim / Co-trimoxazole:</b> Use with caution, may increase K<sup>+</sup> and decrease renal function. Monitor U+Es. If oral route compromised, co-trimoxazole can be given IV (see BNF for dose).</li> </ul>					
Actual Body Weight	Gentamicin Dose	Actual Body Weight	Gentamicin Dose																				
< 40 kg	5 mg/kg	60 - 69 kg	320 mg																				
40 - 49 kg	240 mg	70 - 79 kg	360 mg																				
50 - 59 kg	280 mg	≥ 80 kg	400 mg																				

**INFECTION SPECIALISTS:** Duty Microbiologist, Infectious Disease (ID) Unit at QUEUH. **FOR FURTHER ADVICE:** Clinical/Antimicrobial Pharmacist, local Respiratory Unit (for RTI) or from GGC Therapeutic Handbook.

Infection Control advice may be given by Duty Microbiologist