



## CLINICAL GUIDELINES

# Permanent Pacemaker insertion/ implantable cardiac devices - Antibiotic Prophylaxis recommendations

A guideline is intended to assist healthcare professionals in the choice of disease-specific treatments.

Clinical judgement should be exercised on the applicability of any guideline, influenced by individual patient characteristics. Clinicians should be mindful of the potential for harmful polypharmacy and increased susceptibility to adverse drug reactions in patients with multiple morbidities or frailty.

If, after discussion with the patient or carer, there are good reasons for not following a guideline, it is good practice to record these and communicate them to others involved in the care of the patient.

<b>Version Number:</b>	7
<b>Does this version include changes to clinical advice:</b>	No
<b>Date Approved:</b>	10 <sup>th</sup> May 2022
<b>Date of Next Review:</b>	31 <sup>st</sup> May 2025
<b>Lead Author:</b>	Ysobel Gourlay
<b>Approval Group:</b>	Antimicrobial Utilisation Committee

### Important Note:

The Intranet version of this document is the only version that is maintained. Any printed copies should therefore be viewed as 'Uncontrolled' and as such, may not necessarily contain the latest updates and amendments.



## NHS Greater Glasgow and Clyde recommendations for antibiotic prophylaxis in permanent pacemaker insertion / implantable cardiac devices

Single dose, IV prophylaxis  $\leq$  60 minutes prior to skin incision/ intervention.

**MRSA:** decolonise prior to procedure as per NHS GGC infection control guidelines.

**CPE carriers:** If identified as a Carbapenamase producing Enterobacteriaceae carriers contact microbiology.

Procedure	Recommended Antibiotic	Prolonged Surgery > 4 or > 8 hours
Insertion of permanent pacemaker / implantable cardiac devices	400 mg <b>Teicoplanin</b> $\leq$ 40 Kg  800mg <b>Teicoplanin</b> >40 Kg  Give 400 -800 mg teicoplanin by slow intravenous injection over 3-5 minutes.	No repeat dosing of teicoplanin.