

# GUIDELINE for the REPAIR of a SECOND DEGREE PERINEAL TEAR



<b>TARGET AUDIENCE</b>	Maternity Wide.
<b>PATIENT GROUP</b>	All patients who have sustained a second degree perineal tear.

## Summary

Perineal or genital trauma caused by either tearing or episiotomy is defined as follows:

- First degree – injury to skin only
- Second degree – injury to the perineal muscles but not the anal sphincter
- Third degree- injury to the perineum involving the anal sphincter complex
  - 3a – less than 50% of the external anal sphincter thickness torn
  - 3b – more than 50% of external anal sphincter thickness torn
  - 3c- internal anal sphincter torn
- Fourth degree – injury to the perineum involving the anal sphincter complex (external and internal anal sphincter) and anal epithelium.

### **This guideline is for Second degree tear repair.**

A skilled health care professional should only undertake repair of the perineum following delivery.<sup>1</sup> However, it is acknowledged that this procedure may be performed by a health care professional, who requires to be supervised until competence is achieved. If genital trauma is identified following birth the woman should be advised of the proposed procedure and verbal consent should always be obtained. The initial assessment should be carried out both sensitively and gently immediately following birth.

The woman should be in a position that ensures adequate visual assessment of the degree of trauma. This position should only be maintained for as long as is necessary for the systematic assessment and repair of sustained trauma.

## ***Repair of a second degree perineal tear***

### **Equipment Required**

- Perineal repair pack and associated sterile drapes
- Suture material – Vicryl Rapide 2/0
- X Ray detectable swabs
- Non-alcohol containing cleaning fluid
- Local anaesthetic – Lidocaine 1% 20mls = 200mgs
- Adequate light source
- Clinical waste bag
- Clear plastic bag for swab, tampon and red tag disposal

Once the woman is positioned appropriately the vulva area should be cleansed and sterile drapes arranged as required. Following an initial swab, sharp and instrument count by the clinician and the assistant, the procedure is then undertaken using an aseptic technique.

It is imperative that all swabs, instruments and sharps are counted at the start of the procedure and no assumptions should be made that the starting count is standard. If any additional items are required during the repair then these must be included in the count, and accounted for at the end of the procedure. This must then be documented in badger by the person performing the repair and countersigned by the other person involved in checking.

A further systematic assessment of the perineum and genital tract should be carried out prior to commencing repair. This should include:

- Ensuring the woman is comfortable and local analgesia is effective
- Further explanation of what the health care professional plans to do and why
- Perform visual assessment of the extent of the perineal trauma to include the structures involved, the apex of the injury and assessment of bleeding
- A gentle rectal examination to assess whether there has been any damage to the external or internal anal sphincter.

<b>Lead Author</b>	Gillian Howie	<b>Date approved</b>	September 2024
<b>Version</b>	5	<b>Review Date</b>	September 2027

## ***Repair of a second degree perineal tear***

### **Performing Perineal Repair**

**If repair is following an operative vaginal delivery prophylactic antibiotics are now recommended as per “Antibiotics after Operative Vaginal Delivery for Prevention of Sepsis” guideline**

**Staff are reminded to use appropriate PPE including face masks and eye shields**

1. Unless an effective epidural or spinal anaesthetic is in place, the perineum should be infiltrated with lidocaine 1%. (Refer to Patient Group Directive for dosage details). Confirm the local anaesthesia is working before commencing suturing.
2. Insert a vaginal tampon **only if necessary**, to provide a clearer view of the area to be sutured. The tampon is inserted into the vagina above the level of the apex of the tear and the tab clipped. It is acknowledged that on occasions the tampon may be insufficient in arresting bleeding and allowing the operator a clear view of the area to be repaired. If on these occasions it is necessary to place a swab within the vagina, both the operator and assistant should take a note of this and the swab accounted for in the final swab count. **Only x-ray detectable swabs must be used.**
3. Identify the apex of the wound. Place the first stitch approximately 0.5cm beyond the apex to allow for haemostasis of any small vessels, which may have retracted beyond this point.
4. Insert the anchoring suture above the apex. Repair the vaginal wall using a continuous stitch with approximately 0.5cm between each bite.
5. Carry out the repair from apex to the introitus, ensuring sutures are not placed in the hymenal remnants.
6. At this point place the needle behind the exit point of the last stitch. Sweep it under the fourchette bringing the suture material out into the perineal muscle.
7. Repair the perineal muscles in one or two layers with the same continuous stitch. It is important to oppose the muscle edges carefully and leave no dead space. Usually three or four stitches are all that is required in any one layer.
8. Using a side-to-side technique when suturing subcutaneous (1/2cm bites), continue until the proximal end of the wound is reached. There will be up to a 1/2cm gap between the skin edges.
9. Sweep the needle behind the fourchette back into the vagina. Pick up a small amount of vaginal tissue to tie off the stitch and cut (the knot is tucked into the vagina to minimise discomfort).

<b>Lead Author</b>	Gillian Howie	<b>Date approved</b>	September 2024
<b>Version</b>	5	<b>Review Date</b>	September 2027

## ***Repair of a second degree perineal tear***

10. Check haemostasis has been achieved. NB – an excessive amount of sutures may well cause severe discomfort in the perineum and beyond. Only carry out the required amount of suturing to achieve haemostasis.
11. Remove the tampon or swab, if used, and account for all swabs and needles.
12. Perform a vaginal examination and a rectal examination to confirm that no stitch has penetrated the rectal mucosa.
13. Remove woman's legs from lithotomy position.
14. Explain the extent of trauma and advise woman re hygiene and pain relief associated with perineal trauma.
15. Two health care professionals must check swabs, instruments and needles. All counts of instruments, swabs and needles must be performed and each person must sign as complete in the relevant documentation on badger. The Swabs, tampon and red tag must be placed into a clear bag which is then placed into an orange clinical waste bag for disposal.
16. Document full perineal procedure within Badger.

Lead Author	Gillian Howie	Date approved	September 2024
Version	5	Review Date	September 2027

## ***Repair of a second degree perineal tear***

### **Procedure to be undertaken in the event of an incomplete swab, sharp or instrument count at the end of the perineal repair – Comply with Standard Operating Procedure - Instrument, Sharps and Accountable Items Count Outside Theatre in the Maternity Setting.**

It is imperative that all swabs, instruments and sharps are counted at the start of the procedure and no assumptions should be made that the starting count is standard.

- In the event of the above occurring the clinician and the assistant must repeat the check taking account of all used and unused items.
- The digital vaginal examination should be repeated following gaining consent and providing a full explanation to the patient.
- No material must be removed from the LDRP room or theatre.
- If there is still a discrepancy in the count, midwives and/or medical staff should inform the unit coordinator and consultant of the incident.
- The incident should be reported through Inphase as soon as possible.
- An X-ray must be arranged and a full explanation given to the patient and her partner.

Lead Author	Gillian Howie	Date approved	September 2024
Version	5	Review Date	September 2027

## ***Repair of a second degree perineal tear***

### **References/Evidence**

1. Sultan, AH., Kamm, MA., Hudson, CH., Obstetric Perineal Trauma: An Audit of Training. *Journal of Obstetrics and Gynaecology* 1995; 15: 19-23
2. McElhinney BR, Glenn DRJ, Dornan G Harper MA. Episiotomy repair, vicryl versus vycryl rapide. *Ulster Med J* 2000; 69: 27-29
3. Kettle C, Johanson RB. Continuous versus interrupted sutures for perineal repair. *Cochrane Database System Review* 2003; CD000947
4. Fleming N Can the suturing method make a difference in postpartum, perineal pain. *J Nurse- Midwifery* 1990; 35: 19-25
5. NHSL SOP Swab, Instrument, Sharps and Accountable Items Count Outside Theatre in the Maternity Setting (2023)
6. NHSL guideline Antibiotics after Operative Vaginal Delivery to prevent Sepsis (2020)

<b>Lead Author</b>	Gillian Howie	<b>Date approved</b>	September 2024
<b>Version</b>	5	<b>Review Date</b>	September 2027

**Repair of a second degree perineal tear**

## Appendices

**1. Governance information for Guidance document**

<b>Lead Author(s):</b>	Gillian Howie
<b>Endorsing Body:</b>	Maternity Clinical Effectiveness Group
<b>Version Number:</b>	5
<b>Approval date</b>	September 2024
<b>Review Date:</b>	September 2027
<b>Responsible Person (if different from lead author)</b>	

<b>CONSULTATION AND DISTRIBUTION RECORD</b>	
<b>Contributing Author / Authors</b>	
<b>Consultation Process / Stakeholders:</b>	Maternity CEG Process
<b>Distribution</b>	All in maternity

<b>Lead Author</b>	Gillian Howie	<b>Date approved</b>	September 2024
<b>Version</b>	5	<b>Review Date</b>	September 2027

## Repair of a second degree perineal tear

CHANGE RECORD			
Date	Lead Author	Change	Version No.
2006	Noreen Kent	Initial version (titles of versions 1-4 was " Repair of perineal Lacerations")	1
2012	Noreen Kent	Update	2
2016	Amanda Kennett	Update	3
2020	Jacqueline Holmes	Update	4
			5

### 2.You can include additional appendices with complimentary information that doesn't fit into the main text of your guideline, but is crucial and supports its understanding.

e.g. supporting documents for implementation of guideline, patient information, specific monitoring requirements for secondary and primary care clinicians, dosing regimen/considerations according to weight and/or creatinine clearance

Lead Author	Gillian Howie	Date approved	September 2024
Version	5	Review Date	September 2027