

BNF Chapter	ADR Medication	F	C	UR	CNS	B	HF	Br	CV	R	H	R I	Hypo	Hyp	SS	ACG
1	H2 Blocker	Y	Y	-	-	-	-	-	-	-	-	-	-	-	-	-
	Laxatives	-	-	-	-	-	-	-	-	-	-	-	Y	-	-	-
	Loperamide	Y	Y	Y	Y	-	-	-	-	-	-	-	-	-	-	-
	Prochlorperazine etc ^a	Y	-	-	Y	-	-	Y	Y	-	-	-	Y	-	-	Y
	Metoclopramide	Y	-	-	Y	-	-	Y	Y	-	-	-	-	-	-	-
2	ACE/ARB	Y	Y	-	-	-	-	-	Y	Y	-	Y	-	Y	-	-
	Thiazide diuretics	Y	Y	-	Y	-	-	-	-	-	-	Y	Y	-	-	-
	Loop diuretics	Y	Y	-	-	-	-	-	Y	-	-	Y	Y	-	-	-
	Amiloride ^f / triamterene	Y	Y	-	Y	Y	-	-	-	Y	-	Y	-	Y	-	Y
	Spirolactone	Y	-	-	Y	-	-	-	-	-	-	Y	-	Y	-	-
	Beta-blocker	Y	Y	-	-	-	Y	Y	-	Y	-	-	-	-	-	-
	CCB (dihydropyridine)	Y	Y	-	Y	-	-	-	Y	Y	-	-	-	-	-	-
	CCB (verapamil/ diltiazem)	Y	Y	-	-	-	Y	Y	Y	-	-	-	-	-	-	-
	Nitrates and nicorandil	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Digoxin	Y	-	-	-	-	-	Y	Y	-	-	-	-	-	-	-
3	Theophylline	-	-	-	-	-	-	-	Y	-	-	-	Y	-	-	-
	Oral steroids	Y	-	-	Y	Y	Y	-	-	-	-	-	Y	-	-	Y
4	Opiates	Y	Y	Y	Y	-	-	-	-	Y	-	-	-	-	Y	-
	Benzodiazepines	Y	-	Y	Y	-	-	-	-	Y	-	-	-	-	-	-
	zSedative antihistamines ^d	Y	Y	Y	Y	-	-	-	-	-	-	-	-	-	-	-
	H1 Blockers	Y	Y	Y	Y	-	-	-	-	-	-	-	-	-	-	-
	Antipsychotics ^e	Y	Y	Y	Y	-	-	-	Y	-	-	-	-	-	-	-
	SSRI and related	-	-	Y	Y	Y	-	-	Y	-	-	-	-	-	Y	-
	TCAs ^c	Y	Y	Y	Y	-	Y	-	Y	-	Y	-	-	-	Y	Y
	MAO inhibitors	-	Y	Y	Y	-	-	Y	Y	-	-	-	-	-	Y	Y
5	Antibiotics/ Antifungals	-	-	-	-	-	Y	-	-	-	-	Y	-	Y	-	-
6	Sulfonylureas, gliptins, glinides	Y	Y	-	-	-	-	-	-	-	Y	-	-	-	-	-
	Pioglitazone	Y	-	-	-	-	Y	-	-	Y	Y	-	-	-	-	-
7	Urinary antispasmodics	Y	Y	Y	Y	-	-	-	Y	-	Y	-	-	-	-	Y
	Dosulepin ^b	Y	Y	Y	Y	-	-	-	Y	-	Y	-	-	-	-	Y
	Alpha blocker	Y	-	-	Y	-	-	-	Y	Y	-	-	-	-	-	-
10	NSAIDs	-	-	-	-	Y	-	-	Y	Y	-	Y	-	-	-	-

Key: F - Falls and fractures; C – Constipation; UR - Urinary retention; CNS – CNS Depression; B – Bleeding; HF - Heart failure; Br – Bradycardia; CV - CV events; R –

Respiratory; H – Hypoglycaemia; RI - Renal injury; Hypo – Hypokalaemia; Hype – Hyperkalaemia; SS - Serotonin syndrome; ACG – Angle Closure Angle Glaucoma

- a. strong anticholinergics are: dimenhydrinate, scopolamine, dicyclomine, hyoscyamine, propantheline;
- b. strong anticholinergics are: tolterodine, oxybutynin, flavoxate;
- c. strong anticholinergics are: amitriptyline, desipramine, doxepine, imipramine, nortriptyline, trimipramine, protriptyline;
- d. strong anticholinergics are: promethazine;
- e. strong anticholinergics are: diphenhydramine, clemastine, chlorphenamine, hydroxyzine.
- f. Amiloride side effect frequency unknown